

# A NOTE ON GAMMA RANDOM VARIABLES AND DIRICHLET SERIES\*

Zbigniew J. Jurek,  
(The University of Wrocław, Wrocław, Poland)

25 March 1999

**ABSTRACT.** We give necessary and sufficient conditions for convergence of series of centered gamma random variables. Those series provide distributions from the Lévy class  $L$  of the selfdecomposable probability distributions. Relations to Dirichlet series and the background driving Lévy processes (BDLP's) are investigated.

## 1. The gamma random variable and its selfdecomposability property.

The well-known *gamma random variables* (rv)  $\gamma_{\alpha,\lambda}$  are defined by their densities

$$(1) \quad \lambda^\alpha / \Gamma(\alpha) x^{\alpha-1} \exp(-\lambda x) 1_{(0,\infty)}(x),$$

where  $\alpha > 0$ ,  $\lambda > 0$  are called *shape* and *scale parameters*, respectively, and  $\Gamma(z)$  is the Euler gamma function  $\Gamma(z) := \int_0^\infty x^{z-1} \exp(-x) dx$ ,  $\Re z > 0$ , satisfying the equation  $z\Gamma(z) = \Gamma(z+1)$ . Of course,  $\gamma_{1,\lambda}$  is the exponential rv. (Note: in the sequel we use the same notation for gamma rv and gamma probability density.) Furthermore, it can be easily seen that

$$(2) \quad \mathbb{E}[\gamma_{\alpha,\lambda}] = \alpha/\lambda; \quad \mathbb{E}[\gamma_{\alpha,\lambda}^2] = \alpha(\alpha+1)/\lambda^2; \quad \gamma_{\alpha,\lambda} \stackrel{d}{=} \lambda^{-1}\gamma_{\alpha,1};$$

---

\*Research supported in part by the grant No. 2P03A02914 from KBN, Warsaw, Poland

and also that its characteristic function is

$$(3) \quad \mathbb{E}[\exp(it\gamma_{\alpha,\lambda})] = (1 - it/\lambda)^{-\alpha} = \exp \left[ \int_0^\infty (e^{itx} - 1) \alpha e^{-\lambda x} / x dx \right], t \in \mathbb{R},$$

where  $\mathbb{E}[\cdot]$  denotes the expected value with respect to probability  $\mathcal{P}$ , and  $\stackrel{d}{=}$  means equality in distribution. From (3) we see that

$$(4) \quad \gamma_{\alpha,\lambda} \text{ is infinitely divisible (ID) and its Lévy spectral measure } M \\ \text{(in the Lévy-Khintchine formula) has density } \alpha e^{-\lambda x} / x 1_{(0,\infty)}(x).$$

Recall also here that the class *ID* of all infinitely divisible distributions is a closed convolution semigroup (in weak topology) and it coincides with the class of all processes  $Y$  with stationary and independent increments starting from 0, with probability 1. If additionally  $Y$  has cadlag paths we refer to it as a *Lévy process*.

For  $0 < c < 1$  let us define

$$\psi_c(t) := (1 - ict)/(1 - it) = c1 + (1 - c)/(1 - it), \quad t \in \mathbb{R},$$

which by (3) is a characteristic function (convex combination of 1 and characteristic function of  $\gamma_{1,1}$ ). More precisely,  $\psi_c$  is *ID* with Lévy spectral measure  $x^{-1}(\exp(-x) - \exp\{-c^{-1}x\})1_{(0,\infty)}(x)dx$ . Consequently, for each  $0 < c < 1$  we have a factorization  $\hat{\gamma}_{\alpha,\lambda}(t) = \hat{\gamma}_{\alpha,\gamma}(ct)\psi_c^\alpha(t/\lambda)$  (the selfdecomposability property). Consequently  $\hat{\gamma}_{\alpha,\lambda}$  belongs to class *L* of *all selfdecomposable characteristic functions* defined as follows

$$(5) \quad \phi \in L \quad \text{iff} \quad \forall(0 < c < 1)\exists(\text{char.f. } \phi_c) \phi(t) = \phi(ct)\phi_c(t).$$

The class *L* is a closed subsemigroup in *ID* and coincides with the limiting distributions of normalized partial sums of independent but not necessarily identically distributed rv's. In terms of random rv's selfdecomposability means that rv  $X$  has probability distribution (or char. f.) in class *L* iff  $X \stackrel{d}{=} cX + X_c$ , for each  $0 < c < 1$ , where  $X_c$  is independent of  $X$ ; cf. [7] Chapter 3. Stable laws are the primary examples of class *L* distributions but it is much larger and includes many classical distributions from mathematical statistics like t-Student, F-Fisher, log-normal, etc.; Cf. [6] for other examples. The fundamental characterization of the class *L* is via *the integral functionals of some Lévy processes*. That fact is crucial for the consideration

in this paper. Namely, for class  $L$  distributions or rv's the following *random integral representation* (RIR) holds true:

$$(6) \quad X \in L \quad \text{iff} \quad X \stackrel{d}{=} \int_0^\infty e^{-s} dY(s),$$

where  $Y$  is a Lévy process unique in distribution with finite logarithmic moment  $\mathbb{E}[\log(1+|Y(1)|)] < \infty$ . One refers to  $Y$  as the BDLP for  $X$  (background driving Lévy process). If  $\phi$  is char. f. of  $X$  and  $\psi$  is char. f. of  $Y(1)$  then (6) equivalently reads as follows

$$(7) \quad \log \phi(t) \in L \quad \text{iff} \quad \log \phi(t) = \int_0^1 \log \psi(tu) u^{-1} du,$$

cf. [5], [6], [7]. For the gamma rv we have that its BDLP  $Y$  is the compound Poisson process  $Y(t) = \sum_{n=1}^{\overline{N}_\alpha(t)} \gamma_{1,\lambda}^{(n)}$ , where  $\gamma_{1,\lambda}^{(n)}$ ,  $n = 1, 2, \dots$  are independent copies of gamma rv  $\gamma_{1,\lambda}$  and  $\overline{N}_\alpha(t)$  is Poisson process with intensity  $\alpha$ ; cf. [5], formula (11) and [6] formula (2.2) (note the misprints!). Consequently  $Y(1)$  has compound Poisson distribution with (finite) Lévy spectral measure  $N$  with density

$$(8) \quad dN(x) = \alpha \lambda \exp(-\lambda x) 1_{(0,\infty)}(x) dx$$

[do not confuse here a Lévy spectral measure  $N$  with Poisson process  $\overline{N}(t)$ !].

All in all we have the following:

**PROPOSITION 0.** *Gamma rv's  $\gamma_{\alpha,\lambda}$  are selfdecomposable without Gaussian part and admit representations:*

$$(9) \quad \hat{\gamma}_{\alpha,\lambda}(t) = (1 - it/\lambda)^{-\alpha} = \exp \left\{ \int_0^\infty (e^{itx} - 1) \alpha e^{-\lambda x} / x dx \right\} \\ = \exp \left\{ \int_0^1 \left[ \int_0^\infty (e^{itux} - 1) \alpha \lambda e^{-\lambda x} dx \right] du / u \right\}.$$

This summarizes the study of a single gamma variable. In the next section we will consider sequences of independent gamma random variables.

## 2. Series of independent gamma rv's.

In the sequel  $\gamma_{\alpha_n, \lambda_n}$  denotes a sequence of *independent* gamma rv's. By (2) they belong to  $L^2(\Omega, \mathcal{F}, \mathcal{P})$ , i.e., to the Hilbert space of square integrable rv's with the usual  $L^2$  norm. Thus  $X_n \rightarrow X$  in  $L^2$  if  $\mathbb{E}[|X_n - X|^2] \rightarrow 0$ . Consequently we get:

**PROPOSITION 1.** *A series  $\sum_{n=1}^{\infty} (\gamma_{\alpha_n, \lambda_n} - \alpha_n/\lambda_n)$  of centered gamma rv's converges in  $L^2(\Omega, \mathcal{F}, \mathcal{P})$  iff  $\sum_{n=1}^{\infty} \alpha_n/\lambda_n^2 < \infty$ .*

*Proof.* From the stochastic independence and (2) we get that

$$(10) \quad \mathbb{E}\left(\sum_{n=k}^m \gamma_{\alpha_n, \lambda_n} - \alpha_n/\lambda_n\right)^2 = \sum_{n=k}^m \alpha_n/\lambda_n^2$$

and therefore the completeness of  $L^2(\Omega, \mathcal{F}, \mathcal{P})$  gives the proof.  $\square$

## PROPOSITION 2.

(1) *If  $\sum_n \alpha_n/\lambda_n^2 < \infty$  then  $\sum_n (\gamma_{\alpha_n, \lambda_n} - \alpha_n/\lambda_n)$  converges in distribution.*

(2) *Let denote  $\lambda_\infty := \lim_{n \rightarrow \infty} \lambda_n$  and assume that  $\sum_n (\gamma_{\alpha_n, \lambda_n} - \alpha_n/\lambda_n)$  converges in distribution. Then  $\sum_n \alpha_n/\lambda_n^2 < \infty$  if  $\lambda_\infty > 0$ ; and  $\sum_n \alpha_n < \infty$  if  $\lambda_\infty = 0$ .*

*Proof.* Since  $L^2$  convergence implies convergence in probability which in turn implies convergence in distribution therefore part (1) follows from part (ii) in Proposition 1. For part (2), let us observe that by Proposition 0, formula (9), if  $S$  is the sum of the series in question, then it has Lévy spectral measure

$$(11) \quad dM(x) = \sum_{n=1}^{\infty} \alpha_n e^{-\lambda_n x} / x 1_{(0, \infty)}(x) dx.$$

(in particular the series converges for all  $x > 0$ ). Since Lévy spectral measures integrate  $x^2$  in each finite neighbourhood of zero, we have

$$(12) \quad \int_{|x| < 1} x^2 dM(x) = \sum_n \alpha_n / \lambda_n^2 \int_0^1 \lambda_n x e^{-\lambda_n x} \lambda_n dx = \sum_n \alpha_n / \lambda_n^2 \int_0^{\lambda_n} u e^{-u} du < \infty.$$

The last integral has a limit 2 when  $\lambda_n \rightarrow \infty$  which allows to conclude the first case in part (2). Similarly, when  $\lambda_n \rightarrow 0$  then the integral divided by  $\lambda_n^2$  tends to 1/2 which justifies the second case. The proof is complete.  $\square$

*REMARK 1.* One may try to prove the previous result by the Kolmogorov Three Series Theorem. However, it leads to series of incomplete gamma functions.

**COROLLARY 1.** *Let  $\alpha_n > 0$ ,  $\lambda_n > 0$  be such that  $\inf_n \lambda_n > 0$ . Then for*

$$S_m := \sum_{n=1}^m (\gamma_{\alpha_n, \lambda_n} - \alpha_n / \lambda_n)$$

*the following are equivalent:*

- (a)  $\sum_n \alpha_n / \lambda_n^2$  converges;
- (b)  $(S_m)$  converges in  $L^2(\Omega, \mathcal{F}, \mathcal{P})$ ;
- (c)  $(S_m)$  converges in distribution;
- (d) the series  $h(x) := \sum_n \alpha_n e^{-\lambda_n x}$  converges for all  $x > 0$ , and the integral  $\int_0^1 x h(x) dx < \infty$ .

*Proof.* First of all note that for the claim in Proposition 2 (ii) it is enough to know that  $(\lambda_n)$  are bounded away from zero. Further, (a) implies (b) by Proposition 1, and (b) always implies (c). If  $S$  is the limit of  $S_m$  in distribution then, by Proposition 0,  $h(x)/x1_{(0, \infty)}(x)$  is the density of the Lévy spectral measure of  $S$ . Thus it integrates  $x^2$  in any finite neighbourhood of zero. Thus (c) implies (d). The integrability condition in (d) via (12) implies (a).  $\square$

**COROLLARY 2.** *For the sequence of partial sums*

$$S_m := \sum_{n=1}^m (\gamma_{\alpha_n, \lambda_n} - \alpha_n / \lambda_n),$$

*of independent and centered gamma rv with  $\inf_n \lambda_n > 0$  the following conditions are equivalent:*

- (1)  $(S_m)$  converges a.s.;

- (2)  $(S_m)$  converges in probability;
- (3)  $(S_m)$  converges in distribution;
- (4)  $(S_m)$  converges in  $L^2(\Omega, \mathcal{F}, \mathcal{P})$
- (5)  $M := \sup_m |S_m| \in L^2(\Omega, \mathcal{F}, \mathcal{P})$
- (6)  $K := \sup_n |\gamma_{\alpha_n, \lambda_n} - \alpha_n / \lambda_n| \in L^2(\Omega, \mathcal{F}, \mathcal{P})$ .

*Proof.* Of course, the first three conditions are equivalent for *any* series with independent summands, due to the Lévy Theorem; cf. for instance [2], Theorem 2.10. Condition (3) is equivalent to (4) by Corollary 1. Finally (4),(5) and (6) are equivalent by Hoffmann-Jørgensen's Theorem; cf. Theorem 2.11 in [2].  $\square$

### 3. Dirichlet series.

In complex analysis and notably in the analytic number theory an important role is played by the so called *generalized Dirichlet series*. These are series of the following form:

$$(13) \quad w(z) := \sum_n a_n e^{-\lambda_n z}, \quad \Re z > s_c,$$

where the coefficients  $a_n$  are complex numbers and the exponents

$$0 < \lambda_1 < \lambda_2 < \dots \rightarrow \infty.$$

Of course, Dirichlet series can be viewed as a generalization of power series and a special case of the Laplace transform. For  $\lambda_n = \log n$  we obtain *ordinary Dirichlet series* and in particular the Riemann zeta function. For our purpose here let us recall that (13) converges in half planes, i.e. there exist  $s_c$  (possible  $\pm\infty$ ) such that (13) converges for all  $z \in \mathbb{C}$  with  $\Re z > s_c$  and diverges for all  $z$  with  $\Re z < s_c$ . The line  $\Re z = s_c$  is referred to as *the abscissa of convergence* of the Dirichlet series. The functions  $w(z)$  are holomorphic in the half plane of convergence. Analogously one defines the line  $\Re z = s_a$  as *the abscissa of absolute convergence*; cf. for more details [1] Chapter 8, [8] Chapter 9 or the original work of Harald Bohr in [3] on Dirichlet series and almost periodic functions.

**COROLLARY 3.** *If the series  $\sum_n |a_n|/\lambda_n^2 < \infty$  converges then the generalized Dirichlet series  $\sum_n a_n e^{-\lambda_n z}$  converges absolutely for  $\Re z > 0$ . If additionally the series  $\sum_n |a_n|$  diverges then  $\lim_{m \rightarrow \infty} (\sum_{n=1}^m |a_n|)^{1/\lambda_m} = 1$ .*

*Proof.* The first part follows from Corollary 1 (a) and (d). The second we get from [1] Theorem 8.3.  $\square$

Of course, we have similar statements for ordinary Dirichlet series, i.e. when  $\lambda_n = \log n$ .

*REMARK 2.* Note that having absolutely converging Dirichlet series  $w(z)$  we have class  $L$  distributions with  $w(x)/x1_{(0,\infty)}(x)$  as the density of the Lévy spectral measure  $M$  provided the integrability condition in Corollary 1(d) is satisfied. Moreover,  $-w'(x)$  is the density of the corresponding measure  $N$  from the BDLP  $Y$  ( $N$  is the Lévy spectral measure of  $Y(1)$ ). Conversely, converging series of centered gamma rv's provide examples of summable Dirichlet series in  $\Re z > 0$ . Cf. [5] for some examples.

**4. Thorin's class  $\mathcal{T}$ .** *Generalized gamma distributions* (called also Thorin's class  $\mathcal{T}$ -distributions) are defined via a formula for the moment generating function; see [4], formula (3.1.1) on page 29. For our purpose here it is enough to quote the following fact:

*$\mathcal{T}$  is the smallest class of distributions on  $\mathbb{R}^+$  that contains (shifted) gamma distributions and is closed with respect to convolutions and weak limits;*

cf.[4],Theorem 3.1.5. Furthermore, the so called *Pick functions* (complex analysis) are the main analytic tool. These are functions  $\psi(z)$  that are analytic in the upper complex half-plane  $\Im z > 0$  and have nonnegative imaginary part there; cf. [4], page 20 for more details. From our point of view we obviously have that:

(14) *converging series of centered gamma variables have distributions in  $\mathcal{T}$ ;*

their Lévy spectral measures have densities given by Dirichlet series of the form (13) or their BDLP  $Y$  are infinite series of compound Poisson processes. To conclude what part of class  $\mathcal{T}$  we get from weak limits of infinite series of centered gamma random variables we need to describe limits of sequences of Dirichlet series of the form (13). One can also ask when densities  $h(x)$ ,  $x > 0$ , of Lévy spectral measures  $M$  (in class  $L$ ) admit expansion in form

of a Dirichlet series (13), with coefficients  $\lambda_n$  not necessarily increasing to infinity. Equivalently, one may ask which holomorphic functions  $f(z)$ , in the half plane  $\Re z > 0$  with  $h(x) > 0$ , for  $x > 0$ , and satisfying the integrability condition of Lévy spectral measures, have representation in terms of Dirichlet series. It seems that the original works of Harald Bohr on almost periodic functions and Dirichlet series are still the best reference when one tries to answer those questions.

## REFERENCES

- [1] T. M. Apostol, *Modular functions and Dirichlet series in number theory*. Springer-Verlag, 1976.
- [2] A. Araujo and E. Giné, *The central limit theorem for real and Banach valued random variables*. John Wiley & Sons, New York, 1980.
- [3] H. Bohr, *Collected Mathematical Works*, vol. 1, Dirichlet series, the Riemann zeta-function, København, 1952.
- [4] L. Bondesson, *Generalized gamma convolutions and related classes of distributions and densities*. Lecture Notes in Statistics, v. 76. Springer-Verlag, 1992.
- [5] Z. J. Jurek, Series of independent exponential random variables. *Proc. 7th Japan-Russia Symposium on Probability Theory and Mathematical Statistics, Tokyo 26-30 July 1995*. Ed. S. Watanabe, M. Fukushima, Yu. V. Prohorov, A. N. Shiryaev. World Scientific, Singapore 1996, pp. 174-182.
- [6] Z. J. Jurek, Selfdecomposability: an exception or a rule? *Annales Universitatis M. Curie-Skłodowska*, Lublin-Polonia, Sectio A, vol. LI, 1997, pp. 93-107.
- [7] Z. J. Jurek and J. D. Mason, *Operator-limit distributions in probability theory*. John Wiley & Sons, New York 1993.
- [8] S. Saks and A. Zygmund, *Analytic functions*, 2nd edition, Monografie Matematyczne, Tom 28, PWN-Polish Scientific Publishers, Warszawa 1965.

Author's address:

Institute of Mathematics,  
The University of Wrocław,  
Pl. Grunwaldzki 2/4,  
50-384 Wrocław, Poland.  
[e-mail: zjjurek@math.uni.wroc.pl]