Joint MaPhySto and QUANTOP Workshop on

Quantum Measurements and Quantum Stochastics

Thursday 7 - Tuesday 12 August, 2003
Department of Mathematical Sciences
University of Aarhus

The lectures take place in Auditorium F, Building 534

Program, abstracts and other information for the workshop participants

This booklet contains the programme for the workshop, the abstracts of the talks, general information and a list of participants.

Contents

1 Program **4**

2 Abstracts of Lectures **6**

François Bardou 7

Marcel Bökstedt 7

Vladimír Bužek 8

Ignacio Cirac 9

Giacomo Mauro D'Ariano 10

Christopher A. Fuchs 10

Richard D. Gill 11

Madalin Guta 11

Masahito Hayashi 12

Alexander S. Holevo 12

Peter E. Jupp 13

Jan-Åke Larsson 13

Elena R. Loubenets 13

Hideo Mabuchi 13

Keiji Matsumoto 14

K.R. Parthasarathy 15

Asher Peres 15

Manuel A. Ballester Sanchez 15

Robert Spekkens 16

Jay Gambetta and H. M. Wiseman 17

Christof Wunderlich 17

Jean-Claude Zambrini 18

3 General Information **19**

3.1 Address information 19

3.2 Information Office 19

3.3 Library 19

3.4 Use of computers 19

3.5 Message Board 20

3.6 Meals 20

4 Social arrangements **20**

4.1 Get-together and tour to the QUANTOP laboratory Thursday 7 August, at 5 pm 20

4.2 Excursion to Ebeltoft, Saturday 9 August, from 1 to 7 pm 20

4.3 Workshop Dinner, Monday 11 August, at 7 pm 21

5 Useful Information **21**

5.1 Lounge 21

5.2 Book store 21

5.3 Photocopies 21

5.4 Pay phones 21

5.5 Banks 21

5.6 Pharmacies 22

5.7 Emergencies, doctors and dentist 22

5.8 Casualty ward 22

5.9 Busses close to the department 22

6 List of participants **23**

7 Map of the Department **31**

1 Program

Thursday, August 7

09.00–10.00 Registration and welcome words

10.00–10.50 Alexander S. Holevo (Steklov Mathemtical Institute, Moscow): Asymptotic estimation theory of the shift parameter of a quantum state

11:00–11:50 Ignacio Cirac (Max-Planck-Institut für Quantenoptik, Garching, Germany): Entanglement of Gaussian states and spin systems

LUNCH

14.30–15.20 Keiji Matsumoto (National Institute of Informatics, Quantum Computation Group, Tokyo): Asymptotic quantum estimation theory and Berry-Uhlmann curvature

COFFEE/TEA

15.30–16.20 Christopher A. Fuchs (Bell Labs, Lucent Technologies, New Jersey): What is the difference between a quantum observer and a weatherman?

16:30–17:00 Madalin Guta (Eurandom, Eindhoven): Sieved maximum likelihood state estimation

17:00–19:00 GET-TOGETHER AND LAB TOUR

Friday, August 8

09.00–09.50 Christof Wunderlich (Institut für Laser-Physik, Universität Hamburg): Quantum measurements with individual ions

COFFEE/TEA

10.10–11.00 Elena R. Loubenets (University of Aarhus): On the quantum probabilistic model

11:10–12:00 François Bardou (IPCMS, Strasbourg): Jumps and heavy tails in three microscopic systems

LUNCH

14.30–15.20 K.R. Parthasarathy (Indian Statistical Institute, New Delhi): Construction of non-stabilizer error-correcting quantum codes

COFFEE/TEA

15.30–16.20 Peter E. Jupp (University of St. Andrews): Some aspects of quantum statistical inference

16.30–17.00 Robert Spekkens (Perimeter Institute for Theoretical Physics, Waterloo): In defense of the view that quantum states are states of knowledge

17:00 Discussion

Saturday, August 9

09.00–09.50 Howard Wiseman (Griffith University): The interpretation of non-Markovian stochastic Schrödinger equations: hidden variables

Coffee/tea

10.10–11.00 Asher Peres (Technion, Haifa): Quantum information and relativity theory

11:10–12:00 Marcel Bökstedt (University of Aarhus): Entanglement and geometric invariant theory

Lunch

13.00–19.00 Excursion

Monday, August 11

09.00–09.50 Viacheslav P. Belavkin (University of Nottingham): Quantum causality, stochastics and solution of the measurement problem

Coffee/tea

10.10–11.00 Giacomo Mauro D'Ariano (Northwestern University, Evanston): Universal and programmable measuring devices, and quantum calibration

11:10–12:00 Hideo Mabuchi (California Institute of Technology, Pasadena): Continuous measurement, quantum metrology, and quantum information

Lunch

14.30–15.20 Masahito Hayashi (Quantum Computation and Information Project, ERATO, JST, Tokyo): Quantum central limit theorem and quantum estimation

Coffee/tea

15.30–16.20 Vladimír Bužek (Research Center for Quantum Information, Bratislava): Quantum state reconstruction and optimal manipulations with quantum information

16.30–17.00 Manuel A. Ballester Sanchez (Utrecht University): Estimation of unitary quantum operations

17:00 Discussion

19:00 Workshop dinner

Tuesday, August 12

09.00–09.50 Jean-Claude Zambrini (University of Lisbon): Euclidean quantum mechanics in the momentum representation

Coffee/tea

10.10–11.00 Richard D. Gill (University of Utrecht): Optimal design of the experiment of the century

11:10–12:00 Jan-Åke Larsson (University of Linköping): Bell inequalities for position measurements

Lunch

2 Abstracts of Lectures

François Bardou

IPCMS, 23 rue du Loess, BP 43, F-67034 Strasbourg Cedex 2, France

Jumps and heavy tails in three microscopic systems

This talk will be an introduction to current physics trends which show the importance of jump processes and long tailed distributions in microscopic physical systems.

On the one hand, the observations of individual microscopic quantum or classical objects such as atoms or molecules reveal flagrantly the *jump* nature of many stochastic processes. These jumps were previously masked by ensemble averaging and did not appear explicitly in the formalism. Motivated by the new experimental context, new ways of treating stochastic processes for microscopic systems are emerging, often based explicitly on jumps.

On the other hand, it is being realized that, due to non linear, or even singular, dependences of relevant physical variables on fluctuating parameters generically create *long tails* in the distribution of the physical quantities. These tails generate anomalous scaling behaviours through the inapplicability of the law of large numbers.

The talk will give an overview of three recent physics developments: effects of mutations of RNA molecules [JaB2003], fluctuations in quantum tunneling [DRB2002, RDB2003] and laser cooling of atoms In the last case, a fruitful connection has been established between physics results [BBA2002] and mathematical theories [BBJ2000, BaB2003].

Selected references:

[JaG2003] L. Jaeger and F. Bardou, in preparation.

[DRB2002] V. Da Costa, M. Romeo and F. Bardou, *Statistical properties of currents flowing through tunnel junctions*, J. Magn. Magn. Mater. **258-259**, 90-95 (2002).

[RDB2003] M. Romeo, V. Da Costa and F. Bardou, *Broad distribution effects in sums of log-normal random variables*, Eur. Phys. J. B **32**, 513-525 (2003).

[BBA2002] F. Bardou, J. P. Bouchaud, A. Aspect and C. Cohen-Tannoudji, Lévy Statistics and Laser Cooling, Cambridge University Press, Cambridge (2002).

[BBJ2000] O. E. Barndorff-Nielsen, F. E. Benth and J. L. Jensen, *Markov jump processes with a singularity*, Adv. Appl. Prob. **32**, 779-799 (2000).

[BaB2003] O. E. Barndorff-Nielsen and F. E. Benth, to be published.

Marcel Bökstedt

Department of Mathematical Sciences, University of Aarhus

Entanglement and geometric invariant theory

Let H_1 and H_2 be two finite dimensional vector spaces over the complex numbers. A separable state on the tensor product is a Hermitian operator on $H_1 \otimes H_2$ that can be written as a positive combination of tensor products of Hermitian operators on H_1 respectively H_2.

Even if the dimensions of the vector spaces H_1 and H_2 are as low as 3, it is not easy to describe the geometry of the space of separable states.

I consider the special case where the dimensions of H_1 and H_2 equals 3. The non-compact but reductive group $G = SL_3 \times SL_3$ acts on the Hermitian forms. This action preserves the space of separable hermitian operators.

I use the methods of geometric invariant theory to study this particular case, and to relate the separable states to the stable points of the action of G.

Vladimír Bužek

Research Center for Quantum Information, Slovak Academy of Sciences, Dúbravská cesta 9, 845 11 Bratislava, Slovakia
Department of Mathematical Physics, National University of Ireland, Maynooth, Co.Kildare, Ireland

Quantum state reconstruction and optimal manipulations with quantum informationd

In my talk I will study how information encoded in quantum states can be reconstructed from incomplete experimental data. In particular:
(1) I will show how the *maximum entropy* (MaxEnt) principle can be efficiently used for an estimation of quantum states (i.e., density operators or Wigner functions) on incomplete observation levels, when just a fraction of system observables are measured (i.e., the mean values of these observables are known from the measurement). In particular, I will analyze in detail how the *MaxEnt* principle can be efficiently applied for a reconstruction of quantum states from incomplete tomographic data. I will present reconstruction of

Wigner functions of a single-mode cavity field based on experimental data from a recent experiment of the group of S.Haroche at the Ecole Normale Superior [P.Bertet et al., *Phys. Rev. Lett.* **89**, 200402 (2002)].
(2) When only a *finite* number of identically prepared systems are measured, then the measured data contain only information about frequencies of appearances of eigenstates of certain observables. I will show that in this case states of quantum systems can be estimated with the help of quantum Bayesian inference. I will analyze the connection between this reconstruction scheme and the reconstruction via the MaxEnt principle in the limit of infinite number of measurements.
(3) I will show that the amount of extractable information from finite ensembles is closely related to the fidelity of manipulations that can be performed on physical systems carrying quantum information. Given the bounds imposed by quantum mechanics (all transformations have to be completely positive maps) optimal quantum machines (e.g. universal NOT gates or quantum cloners) can be 'designed'.

Ignacio Cirac

Max-Planck Institut für Quantenoptik, D-85748 Garching, Germany

Entanglement of Gaussian states and spin systems

The entanglement of formation of two particle Gaussian states can be determined by calculating the variance of two specific operators. This quantitiy can be used to determine which are the multipartite Gaussian state, which is invariant under a particular symmetry group, and for which the entanglement of two particular subsystems is maximized. The same question can be asked for qubit systems. In this presentation I will show how to solve these problems and their relation to the entanglement of certain quantum statistical models.

Giacomo Mauro D'Ariano

Department of Electrical and Computer Engineering, Northwestern University,
Evanston, IL 60208

Universal and programmable measuring devices, and quantum calibration

A universal detector acts on a composite quantum system involving an ancilla, and provides the expectation values of any arbitrary operator by just changing the data processing of the outcomes. A programmable detector instead can be programmed to achieve any desired POVM by changing the state of an ancilla. The resources needed to achieve these new kinds of devices are analyzed and quantified, in term of universal machines such as Bell POVM's and control-unitary transformations. Finally a method for giving a quantum calibration of any measuring device with two tomographers and a faithful state is given, with application to a first absolute quantum calibration of a photo-detector.

Christopher A. Fuchs

Bell Labs, Lucent Technologies, Murray Hill, New Jersey 07974

What is the difference between a quantum observer and a weatherman?

Not much. But where there is a difference, there lies quantum theory's most direct statement about properties of the world by itself (i.e., the world without observers or weathermen). In this talk, I will try to make sense of the last two sentences by writing quantum mechanics in terms that only make reference to probability simplexes, rather than Hilbert spaces.

Richard D. Gill

Utrecht University

Quantifying nonlocality-of-entanglement.

One of the uses of entangled states is to do Bell-type experiments. Such an experiment generates data which is incompatible with a classical ("local realist") physical description of the phenomenon. Many different experiments have been proposed with this aim, and some have also been performed: Bell, Clauser-Horne-Shimony-Holt, Mermin, Hardy, Greenberger-Horne-Zeilinger, Popescu, Acin-Gisin-Latorre, ...

Different experiments will need different numbers of particles in order to get equally convincing evidence against local realism. I will quantify this idea, and show how it leads to a new measure of entanglement, or perhaps better, of non-locality: it turns out that the most non-local behaviour is not always observed with the most entangled state.

Reference: quant-ph/0307125 (Van Dam, Gill and Grünwald)

Madalin Guta

Eurandom, Eindhoven

Sieved maximum likelihood state estimation

We describe quantum tomography as an inverse statistical problem and show how entropy methods can be used to study the behaviour of kernel and sieved maximum likelihood estimators.

Masahito Hayashi

Quantum Computation and Information Project, ERATO, JST, Tokyo

Quantum central limit theorem and quantum estimation

Holevo derived a lower bound of mean square error of locally unbiased estimator in quantum estimation. This minimum mean squre error equals to the asymptotic bound of proper estimators. In quantum Gaussian case, the bound by Holevo is achievable. On the other hand, as is proven in quantum central limit theorem, any quantum state goes to quantum Gaussian state. Using this result, I construct an estimator achieving Holevo's bound.

Alexander S. Holevo

Steklov Mathematical Institute, 119991 Moscow

Asymptotic estimation theory of the shift parameter of a quantum state

We develop the estimation theory for the shift parameter in a series of independent copies of a pure quantum state. The optimal covariant estimator is found and its asymptotics is studied. It is shown that under the regularity assumption of finiteness of the second moment of the shift generator, the optimal estimator is asymptotically normal and its variance tends to the absolute lower bound given by the generalized uncertainty relation. On the other hand, the variance of estimators based on unentangled measurements is lowerbounded by a Cramer-Rao bound which can be strictly greater than the bound of uncertainty relation, which is a manifestation of the strict superadditivity of the statistical information in observations of independent quantum systems.

Peter E. Jupp

University of St. Andrews

Some aspects of quantum statistical inference

In classical parametric statistical inference, an important question is 'What parts of the data are informative about the parameters of interest?'. Key concepts here are those of sufficient statistic, ancillary statistic and cut. This talk will outline some analogous concepts for quantum instruments.

Jan-Åke Larsson

Linköping University

Bell inequalities for position measurements

Bell inequalities for position measurements are derived using the bits of the binary expansion of position-measurement results. Violations of these inequalities are obtained from the output state of the nondegenerate optical parametric amplifier.

Elena R. Loubenets

Department of Mathematical Sciences, University of Aarhus

On the quantum probabilistic model

We develop a new mathematical framework for the probabilistic description of an experiment upon a system of any type in terms of initial information representing this system. Based on this framework, we introduce the probabilistic model for the description of non-destructive experiments upon a quantum system and prove that positive bounded linear mappings on the Banach space of trace class operators, arising in the description of generalized quantum measurements, are completely positive.

Hideo Mabuchi

California Institute of Technology, Pasadena

Continuous measurement, quantum metrology, and quantum information

Recent experimental and theoretical work in our group has focused on the use of continuous measurement and real-time feedback to realize and to exploit novel evolutions for optical and atomic systems. In one experiment we used broadband quantum-noise limited measurement and prompt feedback to implement an "adaptive homodyne measurement" that enables single-shot estimation of the phases of weak coherent-state optical pulses, with lower variance than any previously demonstrated technique. In a second, ongoing experiment, we are investigating continuous measurement of the collective spin of a cold atom ensemble. Here, our interests are in spin squeezing, Heisenberg-limited magnetometry, and dynamic generation of large-scale entanglement. In this talk I will review our recent results and discuss the central role of quantum trajectory theory in formulating optimal strategies for metrology and entanglement-generation via continuous quantum measurement.

Keiji Matsumoto

National Institute of Informatics, Quantum Computation Group, Tokyo 101-8430

Asymptotic Quantum Estimation Theory and Berry-Uhlmann curvature

Recently, asymptotic theory of quantum statistics are studied by several authors, and for some cases, optimal efficiency is known. In this talk, pure state models are mainly stdudied, and relation between asymptotic efficiency and Berry phase (Berry-Uhlmann curvature) is discussed. Mixed state models are also studied.

K.R. Parthasarathy

Indian Statistical Institute, New Delhi

Construction of non-stabilizer error-correcting quantum codes

Starting from the well-known theory of classical codes we shall present a brief introduction to the theory of error-correcting quantum codes with emphasis on the Gottesman stabilizer codes. By pooling several Gottesman codes with different character coefficients we shall construct examples of nonstabilizer codes. The connection between the codes of classical information theory and quantum codes on the one hand and the theory of entanglement of states on the other will also be highlighted.

Asher Peres

Department of Physics, Technion - Israel Institute of Technology, Haifa

Quantum Information and Relativity Theory

Quantum mechanics, information theory, and relativity theory are the basic foundations of theoretical physics. The acquisition of information from a quantum system is the interface of classical and quantum physics. Special relativity imposes severe restrictions on the transfer of information between distant systems. Various applications will be presented.

Manuel A. Ballester Sanchez

Utrecht University

Estimation of unitary quantum operation

The problem of optimally estimating an unknown unitary quantum operation is treated. A comparison is made between separable and non-separable measurements and it is shown that non-separable measurements are always better by at least a factor of two. It is also shown that if separable measurements are used, classical communication does not improve

the estimation. In two dimensions it is also shown that measuring one of the Bell states against the other three is as good as any separable measurement and has the advantage of having only two outcomes.

Reference: quant-ph/0305104

Robert Spekkens

Perimeter Institute for Theoretical Physics, Waterloo

In defense of the view that quantum states are states of knowledge

I present a toy theory that is based on a simple information-theoretic principle, namely, that when one has maximal knowledge of a system, one's knowledge is quantitatively equal to one's ignorance. The object analogous to the quantum state in the theory is a uniform probability distribution over hidden states. Because the theory is, by construction, local and non-contextual, it does not reproduce quantum theory. Nonetheless, a wide variety of quantum phenomena have analogues within the toy theory that admit simple and intuitive explanations. Such phenomena include: the non-commutativity of measurements, interference, no information gain without disturbance, the multiplicity of pure state decompositions of a mixed state, the distinction between two-way and intrinsic three-way entanglement, the monogamy of pure entanglement, no cloning, no broadcasting, the remote steering of states, teleportation, dense coding, mutually unbiased bases, locally immeasurable product bases, unextendible product bases, the possibility of key distribution, the impossibility of bit commitment, and many others. The diversity and quality of these analogies provides compelling evidence for the view that quantum states are states of knowledge rather than states of reality, and that maximal knowledge is incomplete knowledge. A consideration of the phenomena that the toy theory fails to reproduce, notably, violations of Bell inequalities and the existence of a Kochen-Specker theorem, provides clues for how to proceed with a research program wherein the quantum state being a state of knowledge is the idea upon which one never compromises.

Jay Gambetta and H. M. Wiseman

Griffith University, Nathan 4111, Australia

The interpretation of non-Markovian stochastic Schrödinger equations: hidden variables

Do diffusive non-Markovian stochastic Schrödinger equations (SSEs) for open quantum systems have a physical interpretation? In a recent paper we investigated this question using the orthodox interpretation of quantum mechanics [Phys. Rev. A **66**, 012108 (2002)]. We found that the solution of a non-Markovian SSE represents the state the system would be in at that time if a measurement was performed on the environment at that time, and yielded a particular result. However, the linking of solutions at different times to make a trajectory is, we concluded, a fiction. In this paper we reinvestigate this question using the modal (hidden variable) interpretation of quantum mechanics. We find that the noise function $z(t)$ appearing in the non-Markovian SSE can be interpreted as the value of a suitably chosen environment property (beable). The solution of the non-Markovian SSE is the state of the system 'conditioned' on this value and the non-Markovian SSE gives its evolution through time. We present the theory for diffusive non-Markovian SSEs that have as their Markovian limit SSEs corresponding to homodyne and heterodyne detection, as well as one which has no Markovian limit.

Christof Wunderlich

Institut für Laser-Physik, Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany

Quantum measurements with individual ions

Individual trapped ions are ideally suited to investigate fundamental issues of quantum mechanics such as the measurement process. Experiments with single Yb^+ ions are reported where an adaptive algorithm for quantum state estimation was implemented, as well as an experiment demonstrating the quantum Zeno paradox with an individual quantum system. In addition, a new type of ion trap is briefly discussed where a collection of ions form an artificial "molecule"' with pairwise spin-spin coupling between individual constituents.

The exact determination of the state of a quantum mechanical two-state system (qubit) would require infinitely many measurements on identically prepared qubits. If a finite

number N of qubits is available, the optimal estimate of the quantum state is obtained by performing a *simultaneous* measurement on all N qubits. We have performed *sequential* measurements on arbitrary but identically prepared states of a single qubit in order to estimate its state more economically. The base of measurement is varied in real time during a sequence of N measurements, conditioned on the results of previous measurements in this sequence.

The quantum Zeno effect is manifest in another experiment that demonstrates a peculiar prediction of quantum mechanics: Even negative-result measurements (that do not involve a local physical interaction in a classical sense) impede the evolution of a quantum system, here the hyperfine states of Yb^+. The analysis of sequences of such null measurements reveals that the quantum system's evolution is indeed reset to "start" (here state F=0) after each negative result measurement.

In a new type of linear ion trap a magnetic field gradient is applied along the axis of rotational symmetry of the trapping potential. In such a trap the internal states (qubits) of different ions are coupled by an effective spin-spin interaction whose formal description is the same as that for J-coupling in molecules used for nuclear magnetic resonance (NMR) experiments. This coupling can be adjusted by choosing the trap parameters and the type of ion. Thus, an artificial ion "molecule´´ can be designed where individual qubits are well isolated from their environment, can be individually addressed, and read out with nearly unity efficiency.

The $S_{1/2}$ ground-state hyperfine doublet (total angular momentum F=0,1) of a single $^{171}Yb^+$ ion confined in a Paul trap represents the two-state system (qubit) in the experiments reported here. The F=0 $\leftrightarrow$ F=1 transition is driven by a quasiresonant microwave (mw) field at 12.6GHz. Resonance fluorescence on the $S_{1/2}$(F=1) $\leftrightarrow$ $P_{1/2}$ transition driven by a laser at 369 nm serves for state-selective detection with efficiency $\eta > 98\%$.

Jean-Claude Zambrini

University of Lisbon

Euclidean Quantum Mechanics in the momentum representation

We give the basis of a (rigorous) probabilistic interpretation of the momentum representation in quantum mechanics. To each diffusion associated with a quantum state in the position representation corresponds a reversible diffusion with jumps driven by Lévy processes in the momentum representation. The strategy used is strongly inspired by Feynman path integral approach, and should be, in fact, independent of the representation.

3 General Information

3.1 Address information

Neetwork in Mathematical Physics and Stochastics
Department of Mathematical Sciences
University of Aarhus
Building 521
Ny Munkegade
DK-8000 Aarhus C
Denmark

MaPhySto Scientific Director Arne Jensen: (+45) 89 42 34 42
Office: A2.34 (building 530 (A), 2nd floor)
MaPhySto Secretary Annemette Hammer: (+45) 89 42 35 32
Office: H2.27 (building 521 (H), 2nd floor)
MaPhySto Secretary Oddbjørg Wethelund: (+45) 89 42 35 15
Office: H2.09 (building 521 (H), 2nd floor)
Fax: (+45) 86 13 17 69

3.2 Information Office

Office hours

Monday to Thursday	8.00 to 15.30
Friday	8.00 to 15.00

3.3 Library

The library is located in building 536 on first, second and third floor.

Opening hours:	8.00 to 15.00
Phone:	89 42 33 67

3.4 Use of computers

The participants may use the computers in the following rooms:

Building 531 (D)	room D02 (ground floor)
Building 531 (D)	room D03 (ground floor)
Building 531 (D)	room D04 (ground floor)
Building 521 (H)	room 2.22 (second floor)
Building 521 (H)	room 3.09 (third floor)

Use login name: `mpsguest`
Password: `Quantum03`
This account will be open only until August 20, 2003.

3.5 Message Board

Please check the message board (placed in front of the auditorium) regularly.

3.6 Meals

Lunch
Lunch is served in the canteen of the Department of Physics and Astronomy, Building 520, 7th floor, for all participants from Thursday to Tuesday, except Sunday. Lunch is free for registered participants and includes one soft drink.

Coffee
During breaks, coffee/tea and cookies will be served outside the auditorium.

Dinner
If you need information on dining in the evenings, please consult the folder "Visit Århus".

4 Social arrangements

4.1 Get-together and tour to the QUANTOP laboratory Thursday 7 August, at 5 pm

4.2 Excursion to Ebeltoft, Saturday 9 August, from 1 to 7 pm

Excursion to Ebeltoft, approx. 1 hour bus transport. Walk in the small, charming town of Ebeltoft, one of the oldest Danish market towns, with possibility to visit the Glasmuseum and Fregatten Jylland. Coffee/tea and cake at the restaurant "Den Skæve Kro", in one of Ebeltoft's oldest buildings. Visit to the Kalø castle ruin. Return to Aarhus at 6 pm.

Please sign up for the excursion (list on the message board) no later than Thursday at 5 pm, thank you.

4.3 Workshop Dinner, Monday 11 August, at 7 pm

5 Useful Information

5.1 Lounge

For the participants and their accompanying persons there is a small lounge in room 2.20, building 530 (A), where there are some international newspapers, magazines and a TV.

5.2 Book store

The Department has its own bookstore located on first floor of building 530 (A). Opening hours are Monday to Friday from 10.00 to 16.00.

5.3 Photocopies

In the Information Office (building 530, first floor) you can buy copy cards for photocopying. A photocopying machine is found in room D04 on the ground floor, close to the canteen.

5.4 Pay phones

Pay phones are available at the entrance to the library on the first floor (across the hall from the main entrance) and in the cloakroom on the ground floor. Phone cards can be bought in the Information Office.

5.5 Banks

Opening hours are Monday, Tuesday, Wednesday and Friday from 10.00 to 16.00.
Thursday: 10.00 to 17.30.
Money machines for credit cards are found outside many banks.

5.6 Pharmacies

Opening hours are Monday to Thursday from 9.30 to 17.30, Friday from 9.30 to 18.00, Saturday from 9.30 to 13.00.
One pharmacy has a 24 hours service. Address:
Løve Apoteket
Store Torv 5, 8000 Århus C. Phone: 86 12 00 22

5.7 Emergencies, doctors and dentist

In case of emergency call **112**.

5.8 Casualty ward

Is found at the "Aarhus Kommunehospital", Nørrebrogade 44, 8000 Århus C and at "Aarhus Amtssygehus", Tage Hansens Gade 2, 8000 Århus C. They are both open 24 hours a day.

The Emergency medical service outside normal hours is available from 16.00 to 8.00 and on weekends available 24 hours a day. Phone: 86 20 10 22.

The Emergency dental services outside normal hours are available from 20.00 to 21.00 all days of the week, and from 9.00 to 10.00 on weekends. Phone 40 51 51 62 will refer you to the dentist on duty.

5.9 Busses close to the department

Busses number 4 (bus stop at Ny Munkegade), 7 and 17 (bus stop at Langelandsgade) go to and from the city centre where they stop right across the street from Hotel Radisson (17), Hotel Ritz (4, 7 and 17) and the railway station.

6 List of participants

Patrik Albin
Department of Mathematical Statistics
Chalmers University of Technology
412 96 Gothenburg
Sweden
Email: palbin@math.chalmers.se

Carlos L. Garrido Alzar
Quantop, Niels Bohr Institute
University of Copenhagen
Blegdamsvej 17
DK-2100 Copenhagen, Denmark
Email: leonardo@nbi.dk

François Bardou
IPCMS
23 rue du Loess
67034 Strasbourg Cedex 2
France
Email: Francois.Bardou@ipcms.u-strasbg.fr

Ole E. Barndorff-Nielsen
Department of Mathematical Sciences
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: oebn@imf.au.dk

Viacheslav P. Belavkin
School of Mathematical Sciences
University of Nottingham
Nottingham NG7 2RD
United Kingdom
Email: viacheslav.belavkin@nottingham.ac.uk

Marcel Bökstedt
Department of Mathematical Sciences
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: marcel@imf.au.dk

Vladimír Bužek
Research Center for Quantum Information
Slovak Academy of Sciences
Dúbrovská cesta 9
845 11 Bratislava, Slovakia
Email: buzek@savba.sk

Ignacio Cirac
Max-Planck-Institut für Quantenoptik
Hans-Lopfermann-Strasse
D-85748 Garching
Germany
Email: ignacio.cirac@mpq.mpg.de

Jeremy Clark
Department of Mathematics
University of California, Davis
One Shields Avenue
95616 Davis, U.S.A.
Email: jeremy@math.ucdavis.edu

Horia Cornean
Department of Mathematical Sciences
Aalborg University
Fredrik Bajers Vej 7E
DK-9220 Aalborg Ø, Denmark
Email: cornean@math.auc.dk

Giacomo Mauro D'Ariano
Quantum Optics and Information Group
Istituto Nazionale di Fisica della Materia
Unita di Pavia, Via A. Bassi 6
I-27100 Pavia, Italy
Email: dariano@unipv.it

Christopher A. Fuchs
Bell Labs
Lucent Technologies
600-700 Mountain Avenue
Murray Hill, NJ 07974, USA
Email: cafuchs@research.bell-labs.com

Richard D. Gill
Mathematical Institute
University of Utrecht
PO Box 80.010
3508 TA Utrecht, The Netherlands
Email: richard.gill@math.uu.nl

Madalin Guta
Eurandom
P.O. Box 513
5600 MB Eindhoven
The Netherlands
Email: guta@eurandom.tue.nl

Peter Harremoës
Department of Mathematics
University of Copenhagen
Universitetsparken 5
2100 Copenhagen, Denmark
Email: moes@math.ku.dk

Masahito Hayashi
Quantum Computation and Information Project, ERATO, JST
5-28-3, Hongo
Bunkyo-ku
Tokyo, 113-0033, Japan
Email: masahito@qci.jst.go.jp

Teiko Heinonen
Department of Mathematics
University of Turku
20014 Turku
Finland
Email: temihe@utu.fi

Ira Herbst
Department of Mathematics
University of Virginia
Kerchoff Hall
22904 Charlottesville, VA, USA
Email: iwh@virginia.edu

Alexander S. Holevo
Steklov Mathematical Institute
Russian Academy of Sciences
117966 Moscow
Russia
Email: holevo@mi.ras.ru

Arne Jensen
Department of Mathematical Sciences
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: matarne@imf.au.dk

Peter E. Jupp
School of Mathematical and Computational Sciences
University of St. Andrews
North Haugh
St. Andrews, KY 16 9SS, Scotland
Email: pej@st-andrews.ac.uk

Jan-Åke Larsson
Matematiska Institutionen
Linköpings University
SE-581 83 Linköping
Sweden
Email: jalar@mai.liu.se

Matthew Leifer
School of Mathematics
University of Bristol
University Walk
BS8 1TW Bristol, United Kingdom
Email: Matt.Leifer@bristol.ac.uk

Elena R. Loubenets
Department of Mathematical Sciences
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: elena@imf.au.dk

Alessandra Luati
Department of Statistics
University of Bologna
Via Belle Arti 41
40126 Bologna, Italy
Email: luati@stat.unibo.it

Hideo Mabuchi
California Institute of Technology
Mail Code 136-93
Pasadena, CA 91125
USA
Email: hmabuchi@caltech.edu

Luis M. Artiles Martinez
Eurandom
P.O. Box 513
5600 MB Eindhoven
The Netherlands
Email: artiles@eurandom.tue.nl

Keiji Matsumoto
National Institute of Informatics
Quantum Computation Group
2-1-2 Hitotsubashi, Chiyoda-ku
Tokyo 101-8430, Japan
Email: keiji@nii.ac.jp

Anders Muszta
Department of Mathematical Statistics
Chalmers University of Technology
412 96 Gothenburg
Sweden
Email: muszta@math.chalmers.se

Klaus Mølmer
Department of Physics and Astronomy
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: moelmer@phys.au.dk

K.R. Parthasarathy
Indian Statistical Institute
S.J.S. Sansanwal Marg
New Delhi-110016
India
Email: krp@isid.isid.ac.in

Asher Peres
Department of Physics
Technion
32000 Haifa
Israel
Email: peres@photon.technion.ac.il

Vivi Petersen
Department of Physics and Astronomy
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: vivip@phys.au.dk

Eugene Polzik
Quantop, Niels Bohr Institute
University of Copenhagen
Blegdamsvej 17
DK-2100 Copenhagen, Denmark
Email: polzik@phys.au.dk

Manuel Ballester Sanchez
Mathematical Institute
University of Utrecht
P.O. Box 80.010
3508 TA Utrecht, The Netherlands
Email: ballester@math.uu.nl

Jürgen Schmiegel
Department of Mathematical Sciences
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: schmiegl@imf.au.dk

Petra Scudo
Technion
Via Roma 78
27050 Torrazza Coste (PV)
Italy
Email: p.scudo@statslab.cam.ac.uk

Erik Skibsted
Department of Mathematical Sciences
University of Aarhus
8000 Aarhus C.
Denmark
Email: skibsted@imf.au.dk

Robert Spekkens
Perimeter Institute for Theoretical Physics
35 King St. N
N2J 2W9 Waterloo
Canada
Email: rspekkens@perimeterinstitute.ca

Raymond Streater
Department of Mathematics
King's College London, Strand
London WC2R 2LS
United Kingdom
Email: rf.streater@kcl.ac.uk

Jens Lykke Sørensen
Department of Physics and Astronomy
University of Aarhus
DK-8000 Aarhus C
Denmark
Email: jls@phys.au.dk

Yonas Tekle
Department of Mathematical Statistics
Chalmers University of Technology
412 96 Gothenburg
Sweden
Email: tekle@math.chalmers.se

Steen Thorbjørnsen
Department of Mathematics and Computer Science
Odense University
Campusvej 55
DK-5230 Odense M, Denmark
Email: steenth@imada.sdu.dk

Howard Wiseman
Centre for Quantum Dynamics
School of Science
Griffith University
Nathan 4111, Australia
Email: h.wiseman@gu.edu.au

Christof Wunderlich
Institut für Laser-Physik
Universität Hamburg
Luruper Chaussee 149
D-22761 Hamburg, Germany
Email: wunderlich@physnet.uni-hamburg.de

Jean-Claude Zambrini
Grupo de Fisica-Matematica
Universidade de Lisboa
Av. Prof. Gama Pinto, 2
1649-003 Lisboa, Portugal
Email: zambrini@alf1.cii.fc.ul.pt

7 Map of the Department



Registration takes place in building 532