

On Integral Equations Arising in the First-Passage Problem for Brownian Motion

GORAN PESKIR*

Abstract

Let $(B_t)_{t \geq 0}$ be a standard Brownian motion started at zero, let $g : (0, \infty) \rightarrow \mathbb{R}$ be a continuous function satisfying $g(0+) \geq 0$, let

$$\tau = \inf \{ t > 0 \mid B_t \geq g(t) \}$$

be the first-passage time of B over g , and let F denote the distribution function of τ . Then the following system of integral equations is satisfied:

$$t^{n/2} H_n \left(\frac{g(t)}{\sqrt{t}} \right) = \int_0^t (t-s)^{n/2} H_n \left(\frac{g(t)-g(s)}{\sqrt{t-s}} \right) F(ds)$$

for $t > 0$ and $n = -1, 0, 1, \dots$, where $H_n(x) = \int_x^\infty H_{n-1}(z) dz$ for $n \geq 0$ and $H_{-1}(x) = \varphi(x) = (1/\sqrt{2\pi}) e^{-x^2/2}$ is the standard normal density. These equations are derived from a single 'master equation' which may be viewed as a Chapman-Kolmogorov equation of Volterra type. The initial idea in the derivation of the master equation goes back to Schrödinger [22].

* Centre for Mathematical Physics and Stochastics (funded by the Danish National Research Foundation).

Mathematics Subject Classification 2000. Primary 60J65, 45D05, 60J60. Secondary 45G15, 45G10, 45Q05, 45K05.

Key words and phrases: The (inverse) first-passage problem, Brownian motion, a curved (non-linear) boundary, a first-passage time, Markov process, the Chapman-Kolmogorov equation, Volterra integral equations (of the first and second kind), a system of nonlinear integral equations.

REFERENCES

- [1] ANDRÉ, D. (1887). Solution directe du problème résolu par M. Bertrand. *C. R. Acad. Sci. Paris* 105 (436-437).
- [2] BACHELIER, L. (1900). Théorie de la spéculation. *Ann. Sci. École Norm. Sup.* 17 (21-86). English translation “*Theory of Speculation*” in “*The Random Character of Stock Market Prices*”, MIT Press, Cambridge, Mass. 1964 (ed. P. H. Cootner) (17-78).
- [3] CHAPMAN, S. (1928). On the Brownian displacements and thermal diffusion of grains suspended in a non-uniform fluid. *Proc. Roy. Soc. London Ser. A* 119 (34-54).
- [4] DOOB, J. L. (1949). Hueristic approach to the Kolmogorov-Smirnov theorems. *Ann. Math. Statist.* 20 (393-403).
- [5] DURBIN, J. (1985). The first-passage density of a continuous Gaussian process to a general boundary. *J. Appl. Probab.* 22 (99-122).
- [6] DURBIN, J. (1992). The first-passage density of the Brownian motion process to a curved boundary (with an appendix by D. Williams). *J. Appl. Probab.* 29 (291-304).
- [7] EINSTEIN, A. (1905). Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen. *Ann. Phys.* 17 (549-560). English translation “*On the motion of small particles suspended in liquids at rest required by the molecular-kinetic theory of heat*” in the book ‘*Einstein’s Miraculous Year*’ by Princeton Univ. Press 1998 (85-98).
- [8] FEREBEE, B. (1982). The tangent approximation to one-sided Brownian exit densities. *Z. Wahrsch. Verw. Gebiete* 61 (309-326).
- [9] FOKKER, A. D. (1914). Die mittlere Energie rotierender elektrischer Dipole im Strahlungsfeld. *Ann. Phys.* 43 (810-820).
- [10] FORTET, R. (1943). Les fonctions aléatoires du type Markoff associées à certaines équations linéaires aux dérivées partielles du type parabolique. *J. Math. Pures Appl.* (9) 22 (177-243).
- [11] HOCHSTADT, H. (1973). *Integral Equations*. John Wiley & Sons.
- [12] ITÔ, K. and MCKEAN, H. P. Jr. (1965). *Diffusion Processes and Their Sample Paths*. Reprint by Springer-Verlag 1996.
- [13] KOLMOGOROV, A. N. (1931). Über die analytischen Methoden in der Wahrscheinlichkeitsrechnung. *Math. Ann.* 104 (415-458). English translation “*On analytical methods in probability theory*” in “*Selected works of A. N. Kolmogorov*” Vol II (ed. A. N. Shiriyayev) Kluwer Acad. Publ. 1992 (62-108).
- [14] KOLMOGOROV, A. N. (1933). Zur Theorie der stetigen zufälligen Prozesse. *Math. Ann.* 108 (149-160). English translation “*On the theory of continuous random processes*” in “*Selected works of A. N. Kolmogorov*” Vol II (ed. A. N. Shiriyayev) Kluwer Acad. Publ. 1992 (156-168).
- [15] LÉVY, P. (1939). Sur certains processus stochastiques homogènes. *Compositio Math.* 7 (283-339).
- [16] MALMQUIST, S. (1954). On certain confidence contours for distribution functions. *Ann. Math. Statist.* 25 (523-533).

- [17] PARK, C. and PARANJAPÉ, S. R. (1974). Probabilities of Wiener paths crossing differentiable curves. *Pacific. J. Math.* 53 (579-583).
- [18] PARK, C. and SCHUURMANN, F. J. (1976). Evaluations of barrier-crossing probabilities of Wiener paths. *J. Appl. Probab.* 13 (267-275).
- [19] PESKIR, G. (2001). Limit at zero of the Brownian first-passage density. *Research Report* No. 420, *Dept. Theoret. Statist. Aarhus* (12 pp).
- [20] PLANCK, M. (1917). Über einen Satz der statistischen Dynamik and seine Erweiterung in der Quantentheorie. *Sitzungsber. Preuß. Akad. Wiss.* (324-341).
- [21] RICCIARDI, L. M., SACERDOTE, L. and SATO, S. (1984). On an integral equation for first-passage-time probability densities. *J. Appl. Probab.* 21 (302-314).
- [22] SCHRÖDINGER, E. (1915). Zur Theorie der Fall- und Steigversuche an Teilchen mit Brownscher Bewegung. *Physik. Zeitschr.* 16 (289-295).
- [23] SIEGERT, A. J. F. (1951). On the first passage time probability problem. *Phys. Rev.* (2) 81 (617-623).
- [24] SMOLUCHOWSKI, M. v. (1913). Einige Beispiele Brown'scher Molekularbewegung unter Einfluss äußerer Kräfte. *Bull. intern. Acad. Sc. Cracovie A* (418-434).
- [25] SMOLUCHOWSKI, M. v. (1915). Notiz über die Berechnung der Brownschen Molekularbewegung bei der Ehrenhaft-Millikanschen Versuchsanordnung. *Physik. Zeitschr.* 16 (318-321).
- [26] STRASSEN, V. (1967). Almost sure behavior of sums of independent random variables and martingales. *Proc. Fifth Berkeley Symp. Math. Statist. Probab.* (Berkeley 1965/66) Vol II, Part 1, *Univ. California Press* (315-343).
- [27] ZUCCA, C. (2001). *Analytical, numerical and Monte Carlo techniques for the study of the first passage times*. PhD Thesis, University of Milano.